

NSDUH 2013-2014 Vermont Data Summary

Prevalence and Quintile Rankings*

| | Past Month % | VT Rank 1-5 | Past Year % | VT Rank |
|---|--------------------|-------------------|-------------------|------------|
| Alcohol (12+) | 61.2 | 1 | | |
| Underage Consumption (12-20) | 29.9 | 1 | | |
| Underage Binge (12-20) | 20.3 | 1 | | |
| 12-17 Consumption | 13.8 | 1 | | |
| 12-17 Binge | 7.5 | 1 | | |
| 18-25 Consumption | 71.5 | 1 | | |
| 18-25 Binge | 46.8 | 1 | | |
| 26+ Consumption | 64.4 | 1 | | |
| 26+ Binge | 21.0 | 4 | | |
| 18+ Consumption | 65.4 | 1 | | |
| 18+ Binge | 24.8 | 3 | | |
| Marijuana (12+) | 13.2 | 1 | 20.0 | 1 |
| 12-17 | 11.4 | 1 | 18.3 | 1 |
| 18-25 | 30.6 | 1 | 46.3 | 1 |
| 26+ | 10.4 | 1 | 15.6 | 1 |
| 18+ | 13.4 | 1 | 20.1 | 1 |
| | | | Cocaine | |
| Illicit Drugs Other than Marijuana (12+) | 3.7 | 1 | 2.3 | 1 |
| 12-17 | 3.3 | 3 | 0.8 | 1 |
| 18-25 | 8.6 | 1 | 7.4 | 1 |
| 26+ | 2.9 | 2 | 1.5 | 2 |
| 18+ | 3.7 | 3 | 2.4 | 1 |
| Non-Medical Use of Pain Relievers – NMUPR (12+) | | | 3.3 | 5 |
| 12-17 | | | 4.0 | 5 |
| 18-25 | | | 7.6 | 4 |
| 26+ | | | 2.5 | 5 |
| 18+ | | | 3.2 | 5 |
| RISK PERCEPTION | | | | |
| Great Risk of Binge Drinking 1,2 Times/Week (12+) | 37.0 | 2 | | |
| 12-17 | 31.2 | 1 | | |
| 18-25 | 24.2 | 1 | | |
| 26+ | 39.8 | 2 | | |
| 18+ | 37.5 | 2 | | |
| Great Health Risk of Smoking Marijuana 1/Month (12+) | 18.6 | 1 | | |
| 12-17 | 18.9 | 1 | | |
| 18-25 | 7.1 | 1 | | |
| 26+ | 20.6 | 1 | | |
| 18+ | 18.6 | 1 | | |

* Rank among 50 States and DC (1 = highest 10, 5 = lowest 10)

Quintiles more accurately reflect rankings due to instability in year-to-year individual ranks

Red in Rankings indicates VT in first quintile (bottom quintile for risk perception)

Note : **Green in prevalence estimates** indicates significant reductions from 2012-2013. **Red** in prevalence estimates indicates a significant increase from 2012-2013 (decreases for perception of risk).